

4-7-2022

## Environmental And Diagnostic Study Of Some Parasites In Barbus Xanthopterus Fish Of Al-Diwaniyah River

Abtisam Jasim Abbas

Follow this and additional works at: <https://qjps.researchcommons.org/home>



Part of the [Environmental Sciences Commons](#)

---

### Recommended Citation

Abbas, Abtisam Jasim (2022) "Environmental And Diagnostic Study Of Some Parasites In Barbus Xanthopterus Fish Of Al-Diwaniyah River," *Al-Qadisiyah Journal of Pure Science*: Vol. 27: No. 1, Article 24.  
DOI: 10.29350/qjps.2022.27.1.1533  
Available at: <https://qjps.researchcommons.org/home/vol27/iss1/24>

This Article is brought to you for free and open access by Al-Qadisiyah Journal of Pure Science. It has been accepted for inclusion in Al-Qadisiyah Journal of Pure Science by an authorized editor of Al-Qadisiyah Journal of Pure Science. For more information, please contact [bassam.alfarhani@qu.edu.iq](mailto:bassam.alfarhani@qu.edu.iq).



## Environmental and diagnostic study of some parasites in *Barbus Xanthopterus* fish of Al-Diwaniyah River

<p><b>Authors Names</b></p> <p>a. Abtisam Jasim Abbas</p> <p><b>Article History</b></p> <p>Received on: 10/4/2022 Revised on: 20/5/2022 Accepted on: 5/6/2022</p> <p><b>Keywords:</b> <i>Barbus Xanthopterus, Chilodonella Cyprinid and Eimiria sinensis.</i></p> <p><b>DOI:</b><a href="https://doi.org/10.29350/jops.2022.27.1.1533">https://doi.org/10.29350/jops.2022.27.1.1533</a></p>	<p><b>ABSTRACT</b></p> <p>The current study was concerned with investigating parasitic infections in <i>Barbus Xanthopterus</i> fish in Al-Diwaniyah River, for the period (July 2020-December 2020). (218) fish were collected from three different locations along the Al-Diwaniyah River. The results of the laboratory examination indicated that the fish were infected with the parasite <i>Chilodonella Cyprinid</i> and <i>Eimiria sinensis</i>. The current study recorded changes in the physical and chemical properties of Al-Diwaniyah River water, as the water temperature values, measured turbidity values, pH, electrical conductivity, hardness and TDS recorded a clear variation during the months of the study. The results of the statistical analysis indicated that there were significant differences between these variables at a level of probability (<math>P \leq 0.05</math>).</p>
---	---

### Introduction

In the environment, fish are exposed to many problems that affect their reproduction and threaten their survival, such as competition, parasitism and predation (1). Fish parasites are one of the most important obstacles to the growth and reproduction of fish, in addition to their negative impact on the economic side, through high mortality in small fish or the growth of fish with low weight (2). In general, parasites cause chemical and mechanical damage to their hosts, negatively affecting various physiological activities such as sun, reproduction and digestion (3). Infection with fish parasites depends on the geographical location of the water body in which the

fish live, the depth of the river, the type of animals and plants present (4). Parasites are affected by the physiological, phenotypic and life characteristics of the host, such as the diet of the host, the age and sex of the host, the time of reproduction and activity (4). Parasites migrate through the connective tissues of fish, which hinders growth and reproduction (5). A number of fish parasites or their larval stages are transmitted to humans as a result of eating raw or undercooked fish (5). The current study aims to investigate some types of parasites in *Barbus Xanthopterus* fish, Al-Diwaniyah River.

### Material and method

*Barbus Xanthopterus* fish samples were collected from fishermen in the Al-Diwaniyah River, with a rate of 3 replicates per month. That was during the study period that lasted six months, July-2020 until December 2020. The number of examined fish reached (218). The fish were transferred to the laboratory within a few hours, the fish were subjected To physical examination by noting the injuries (scars, bleeding, falling scales, ulcers, color and quantity of mucous surrounding the body of the fish). DPX, then the slide was loaded according to the method (6). Parasites were diagnosed by taking measurements of each parasite and drawing it with a reconnaissance camera, then diagnosis according to (7).

The Water samples were taken in three locations in the Al-Diwaniyah River, two samples per month during the study period, using opaque plastic bottles. The samples were taken to the laboratory for physical and chemical analyzes, in addition to field experiments that included: Water temperature using [a mercury thermometer], Turbidity measure by [the Turbidity meter type HANA Lp 2000], EC measure by using [Electrical Conductivity type H99300 HANNA, PH using electronic PH meter type Iovibond 150 Senso Direct.

### Results and discussion

(218) *Barbus Xanthopterus* fish were collected from three areas along Al-Diwaniyah River in Al-Diwaniyah Governorate and examined for parasitic infestations. The results indicated that the total infection rate was 39.44%, or 86 samples out of 218. Table (1):

**Table (1): the number of fish samples and the total infestation rate during the months of the study.**

months of study	No. of fish samples	No.of infected cases	Total infection rate
-----------------	---------------------	----------------------	----------------------

	examined		
July	39	18	%46.15
August	42	19	%45.23
September	49	22	%44.89
October	32	13	%40.62
November	30	10	%33.33
December	26	4	%15.38
Total	218	97	%44.49

By investigating the type of parasite, the results indicated that *Barbus Xanthopterus* was infected with the primary parasite *Chilodonella Cyprinid* (moroff, 1902).

**This parasite settles in the skin and gills of infected fish. It belongs to:**

Phyllum: Ciliophora

Class: Cilita

Order: Phllopharyngia

Family: Chlamydodontidae

**The results indicated that the species *Eimiria sinensis*, which belongs to:**

Class: Sporozoa

Order:Coccidia

Family:Emeridae

The parasite infection rates were recorded for the two species (*Chilodonella Cyprinid*, *Eimiria sinensis*) (56.7%-43.29%), respectively. Table (2)

**Table (2): Infection rate of different genera and identified species in *Barbus Xanthopterus* fish.**

Diagnosed parasites	No.of samples of infected fish	%Infection rate
<i>C.cyprinid</i>	55	%56.7

<b>E.sinensis</b>	<b>42</b>	<b>743.29</b>
-------------------	-----------	---------------

### Physical and chemical properties of Al-Diwaniyah River

Observing the characteristics of the Al-Diwaniyah River is one of the most important environmental studies, especially the physical and chemical properties of the river. The living organisms affected by seasonal and locational variations and the ecological condition of the river, table (4):

**Table (3) Arithmetic mean, standard deviation and range (first grade), upper and lower values of physical and chemical Parameters (second grade) in Al-diwanayah River during stations**

Parameters	Stations		
	1	2	3
Water Temp.	16.314±5.261a	16.008±5.400 a	17.030±5.216 a
	8-26	10.87-17	10-27
pH	7.663±0.245 a	7.642±0.130 a	7.710±0.115 a
	7.2-8.2	7.31-7.9	7.5-7
T.D.S	851.418±146.644 a	872.285±130.418 a	876.429±158.400 a
	630-1011	660-1011	650-1112
Turbidity	8.313±2.618 <b>b</b>	13.560±6.213 <b>a</b>	13.475±6.630 <b>a</b>
	3-12	8-26.7	8-16.7
D.O	7.645±1.042 a	7.518±0.984 a	7.540±1.0320 a
	7-9	6-9.5	6-8.7
Tot-Hardness	511.143±8.552 <b>a</b>	480.618±36.750 <b>b</b>	451.275±56.642 <b>c</b>
	505-523	430-517	360-515

The averages carrying different letters within the same row differ significantly between each other

**Table (4): Arithmetic mean, standard deviation and range of parameters of the study period from July-2020 to December-2020**

Parameters	July 2020	Ogeust	September	October	November	December
	Water Temp.	25±1.2 a 23±26	21.064±1.320 b 20±23	12.023±1.140 d 11±14	10.877±0.783 e 9±11	13.123±0.608 d 12±13
pH	7.526±0.074 c 7.32-7.63	7.6766±0.158 bc 7.3-8	7.732±0.147 ab 7.4-8	7.56±0.193 a 7.6-8.2	7.766±0.170 ab 7.2-8	7.622±0.131 bc 7.3-7.3
T.D.S	988±19.025 b 960-1022	1007±14.281 ab 970-1013	1038.656±51.253 a 980-1111	926.6656±14.68 c 900-940	800±71.926 d 710-900	674±11.510 e 650-6958
Turbidity	22.266±7.711 a 11-27.7	16.422±4.032 b 11-20	9.332±2.4 cd 6-13	12±3.112 bc 8-15	10.156±1.447 cd 7-11	7.6656±2.168 d 3-9
D.O	6.322±0.210 d 6-5.6	7.066±0.710c 5-7	7.556±0.370 b 7-6	9.3±0.364 a 8-9	7.956±0.516 b 6-8	7.7±0.224 b 7.4-8.0
Tot-Hardness	504.767± 11.934 a 510-470 0.54-0.7	510±7.4 a 520-500 0.6-0.8	513.322±7.286 a 514-500 0.7-1.5	517.322±9.300 a 518-500 1.27-1	482.656±48.548 a 523-410 0.5-1.2	447.322±64.951 b 521-360 0.5-0.7

The current study indicated that the water temperature varies with the different months of the year. Water temperature is an important determinant of water density, which makes it a determining factor in the distribution of organisms in water bodies (8). This factor has an important role in the photosynthesis of aquatic plants and the decomposition of organic matter, which affects the pH, dissolved oxygen and dissolved carbon dioxide gas (9,10, 11).

The results indicated that the pH values corresponded to the basal values. The baseline is affected by temperature, the increase in the decomposition of organic matter, the increase in the concentration of (Co<sub>2</sub>), the high water levels and the concentration of magnesium (12). The reason for the decrease in PH in some hot months, is the consumption of free Co<sub>2</sub> by the producing organisms, in addition to the presence of carbonate ions, bicarbonates and dissolved silicate salts in water (13).

The results of the electrical conductivity values, as well as the values of dissolved oxygen and the values of the vital oxygen requirement, varied during the months of the study. The results showed varying levels of turbidity. The reason for this result is due to the high speed of water flow and the mixing and raising of water deposits by water currents (9). The increase in turbidity may be due to low water levels in some months and high temperatures that result in increased rates of evaporation, and the fluctuation of the speed of its flow changes the turbidity value of the water as a result of suspended substances carried by the river when it flows (14,15). The rise in hardness in some months is due to the high values of calcium carbonate concentrations and suspended and dissolved substances in the river (16), while (17) indicated that its rise results from a decrease in the water level, which results in an increase in the concentration of ions causing water hardness. In general, it increases in the summer months as a result of dust storms that are loaded with dust particles, which are a source of calcium carbonate (18).

The results indicate the high values of TDS concentrations in the Al-Diwaniyah River. The reason for this increase may be the decrease in the river water level or the speed of water flow during the time of sampling and measuring the parameters under study. As the speed of the flow and the decrease in the water level is a reason for the increase in the concentration of the components and the increase in the concentration of salts. (14,15).

## References

- 1-Nikolsky, G. V. (1963). "The ecology of fishes (Engl. translation)", Acad. Press, London & New York: 352 pp.
- 2-Al-Nari, A. A. W. (1997). "Study on history changes in gonads of *Acanthobrama* named infected with plerocercoid of *Ligula intestinalis*", M. Sc. Thesis, College of Sciences, University of Mosul, Iraq.
- 3-Khalifa Ahmed Khalifa, (1986). *Fish Diseases*", second edition, Directorates of Dar Al-Kutub for Printing and Publishing, University of Mosul, p. 266.
- 4- Dogiel, V. A. , (1961). Ecology of the parasites of freshwater fishes. In: Dogiel, V. A.; Petrushevski, G. K. & Polyanski, Yu. I. (eds.). *Parasitology of fishes (English translation)*", Oliver and Boyd, Edinburgh and London: 1-47.
- 5-Hoffmann, G. L. (1999). *Parasites of North American freshwater fishes*. 2nd ed. Comstock publishing Associates, Ithaca, New York.
- 6- Scholz, T. and Aguirre –Macedo, M. L. (2000). "Metacercariae of trematodes parasitizing freshwater fish in Mexico: a reappraisal and methods of study. In: Salgado – Maldonado, G.; Garcia – Aldrete, A. N. and Vidal- Martinez, V. M. (eds. )", *Metazoan parasites in the Meotrops: a systematic and ecological perspective*, Inst. Piol., Univ. Nac. Auton. Mexico: 101- 115.

- 7-Yamaguti, S. (1961). "The nematodes of vertebrates, part I+II. Intersci", Systema helminthun. Vol. III, The nematodes of vertebrates, part I+II. Intersci, Publ., New York, 1261,
- 8- Smith, R. (2004). " Current methods in aquatic science ", University of Waterloo, Canada.
- 9- Flowers of Younis Reda Muhammad Al-Assaf, (2009) .Wadi Al-Dangali in the city of Mosul", Master's thesis / College of Education - University of Mosul.
- 10- Al-Khalidy, K. A. H.; Al-abodi, H. R. J. and almaehi, A. M. Y. (2018). Investigation of the biological pollution in Al-Diwaniyah River (branch of Euphrates River) in Al-Qadisiyah province. Iraq. *International Journal of Environment & Water*; 7( 3): 92-101.
- 11- Hussein, S. A. Essa, S. A. and Al-Manshed, (2000).Limnological investigations to the lower reaches of Saddam River. 1.Environmental characterisic . Basrah ., *J.Agric. Sci.* 13 (2).
- 12-Hussian, S. A., AL-sabonji, A. A. and Fahad, K. K. (2008). Environmental characteristics of the Euphrates River in the city of Nasiriyah, seasonal differences in some physical and chemical factors . *Dhi Qar University Journal.* (1) (4): 20-13.
- 13- Alkhalidi, S. K. and Al-abodi, H. A. (2021). BIOLOGICAL AND ENVIRONMENTAL STUDY OF AL-DIWANIYAH RIVER – IRAQ. *Turkish Journal of Physiotherapy and Rehabilitation*; 32(3): 11453-11458. <https://turkjphysiotherrehabil.org/pub/pdf/321/32-1-1438.pdf>
- 14- Zidan, T. A., Abdul-Rahman, A. A. and Suud, O. M. (2009). Study of the environment of chemical and physical pollutants affecting the waters of the Euphrates River in Ramadi and Fallujah. *Anbar University Journal of Pure Sciences.* 3(3).
- 15- Zerine, T., Islam, A., Gulnazar, S., Farjana, N. E., Begum, M. A. ., & Sadia, H.-E. (2021). Identification and Antibiotic Susceptibility of Blood Culture Isolates from Rajshahi, Bangladesh. *Journal of Scientific Research in Medical and Biological Sciences*, 2(2), 1-10. <https://doi.org/10.47631/jsrmb.v2i2.264>
- 16-Mohamed, A. J. (2012). Study of the water environment of Yusufiya River, south of Baghdad. *Tikrit Journal of Science and Pure.* 17 (3): 38-48.
- 17-Salman, J. M. and Saud, A. A. (2015). A study of the variability of some environmental factors and evidence of biological diversity of aquatic plants in the Hilla River in central Iraq. *Babylon University Journal of Pure and Applied Sciences.* 1 (35): 114-166.
- 18-Halbeisen, J. (2003) Rely on pH for calcium needs. *The Growers Solutions*, vol. 16, issue 4, p: 1-4

