

4-7-2021

Faber Polynomial for Holomorphic Functions Involving Differential Operator

Huda Fawzi Hussian

Department of Mathematics, College of Education for pure science, University of Anbar-Iraq,
huda.fawzi@uoanbar.edu.iq

Abdul Rahman Salman Juma

Department of Mathematics, College of Education for pure science, University of Anbar-Iraq,
dr_juma@hotmail.com

Follow this and additional works at: <https://qjps.researchcommons.org/home>



Part of the [Mathematics Commons](#)

Recommended Citation

Hussian, Huda Fawzi and Juma, Abdul Rahman Salman (2021) "Faber Polynomial for Holomorphic Functions Involving Differential Operator," *Al-Qadisiyah Journal of Pure Science*: Vol. 26: No. 2, Article 6.
DOI: 10.29350/qjps.2021.26.2.1294

Available at: <https://qjps.researchcommons.org/home/vol26/iss2/6>

This Article is brought to you for free and open access by Al-Qadisiyah Journal of Pure Science. It has been accepted for inclusion in Al-Qadisiyah Journal of Pure Science by an authorized editor of Al-Qadisiyah Journal of Pure Science. For more information, please contact bassam.alfarhani@qu.edu.iq.



Faber Polynomial for Holomorphic Functions Involving Differential Operator

<p>Authors Names a. Huda Fawzi Hussian b. Abdul Rahman Salman Juma</p> <p>Article History Received on: 27/2/2021 Revised on: 25/3/2021 Accepted on: 28/3/2021</p> <p>Keywords: Univalent function, Holomorphic function, Faber polynomial, Differential operator.</p> <p>DOI:https://doi.org/10.29350/jops.2021.26.2.1294</p>	<p>ABSTRACT</p> <p>The purpose of this paper, is to present differential operator for the univalent functions employ a Faber Polynomial . In addition, we will introduce some inclusion properties of the operator that were obtained employ the principle of subordination between holomorphic functions.</p>
---	---

1.Introduction

Let \mathcal{A} be the class of functions f of the form

$$f(z) = z + \sum_{k=2}^{\infty} a_k z^k, (z \in \mathcal{U}) \quad (1.1)$$

which are holomorphic and normalized in the unit disk $\mathcal{U} = \{z \in \mathbb{C} : |z| < 1\}$.

Definition:1.1. Let $g(z)$ be holomorphic and univalent in \mathcal{U} and f also holomorphic in \mathcal{U} , then f is subordinate to g if there be a Schwartz functions $w(z)$ it is also holomorphic in \mathcal{U} with $w(0)=0$ and $|w(z)| < 1, \forall z \in \mathcal{U}$ such that $f(z) = g(w(z))$.

This denoted by $f \prec g$. Furthermore, assume g is univalent in \mathcal{U} . Then the next part holds [4,5,8,10]:

$$f \prec g \Leftrightarrow f(0) = g(0) \text{ and } f(\mathcal{U}) \subset g(\mathcal{U}).$$

For the function f of the form (1.1), we will present result the following: f is starlike and convex, respectively, with respect to the origin, if and only if

$$\operatorname{Re} \left\{ \frac{zf'(z)}{f(z)} \right\} > 0, \quad |z| < 1,$$

and

$$\operatorname{Re} \left\{ 1 + \frac{zf''(z)}{f'(z)} \right\} > 0, \quad |z| < 1,$$

f is starlike and convex, respectively, of order τ , if only and if

$$\operatorname{Re} \left\{ \frac{zf'(z)}{f(z)} \right\} > \tau, \quad 0 \leq \tau < 1, \quad |z| < 1,$$

and

$$\operatorname{Re} \left\{ 1 + \frac{zf''(z)}{f'(z)} \right\} > \tau, \quad 0 \leq \tau < 1, \quad |z| < 1.$$

Remark 1.1 The function f is convex, if and only if $zf'(z)$ is starlike. This is according to the Alexander's Theorem [3].

Definition: 1.2. Let $f \in \mathcal{A}$ and g is starlike of order τ , i.e. $g \in S^*(\tau)$, then $f \in K(\varepsilon, \tau)$, if and only if, $\operatorname{Re} \left\{ \frac{zf'(z)}{g(z)} \right\} > \varepsilon, z \in \mathcal{U}$, we named this function closed-to-convex function of order ε , type τ . For $f \in \mathcal{A}$, the next subclasses of starlike, convex and closed-to-convex functions $S^*(\zeta, \varpi)$, $C(\zeta, \varpi)$ and $K(\zeta, \rho; \varpi, \varphi)$ of order ζ , with many authors have studied [7,8,9,10], and respectively defined by:

$$S^*(\eta, \vartheta) = \left\{ f \in \mathcal{A} : \frac{1}{1-\eta} \left(\frac{zf'(z)}{f(z)} - \eta \right) \prec \vartheta(z), z \in \mathcal{U} \right\}$$

$$C(\eta, \vartheta) = \left\{ f \in \mathcal{A} : \frac{1}{1-\eta} \left(1 + \frac{zf''(z)}{f'(z)} - \eta \right) \prec \vartheta(z), z \in \mathcal{U} \right\}$$

$$K(\eta, \xi; \vartheta, \varphi) = \left\{ f \in \mathcal{A} : \frac{1}{1-\xi} \left(\frac{zf'(z)}{g(z)} - \xi \right) \prec \varphi(z), z \in \mathcal{U}, g(z) \in S^*(\eta, \vartheta) \right\}$$

From (1.1), we have

$$f(z)^\alpha = z^\alpha \left[1 + \alpha(a_2z + a_3z^2 + a_4z^3 + \dots) + \frac{\alpha(\alpha-1)}{2!} (a_2z + a_3z^2 + a_4z^3 + \dots)^2 + \dots \right]$$

$$\begin{aligned}
&= z^\alpha + \alpha a_2 z^{\alpha+1} + \left(\alpha a_3 + \frac{\alpha(\alpha-1)}{2!} a_2^2 \right) z^{\alpha+2} + \left(\alpha a_4 + \frac{\alpha(\alpha-1)}{2!} 2a_2 a_3 + \frac{\alpha(\alpha-1)(\alpha-2)}{3!} \right) z^{\alpha+3} \\
&\quad + \left(\alpha a_5 + \frac{\alpha(\alpha-1)}{2!} (2a_2 a_4 + a_3^2) + \frac{\alpha(\alpha-1)(\alpha-2)}{3!} 3a_2^2 a_3 + 6 + \frac{\alpha(\alpha-1)(\alpha-2)(\alpha-3)}{4!} \right) z^{\alpha+4} , \\
&= z^\alpha + \alpha a_2 z^{\alpha+1} + \alpha a_3 z^{\alpha+2} + \alpha a_4 z^{\alpha+3} + \dots
\end{aligned}$$

definition we give $(f(z))^\alpha$ by

$$f^\alpha(z) = z^\alpha + \sum_{k=2}^{\infty} a_k(\alpha) z^{\alpha+k-1}. \quad (1.2)$$

Thus as $\alpha \rightarrow 1$, to have :

$$f(z) = z + \sum_{k=2}^{\infty} a_k z^k.$$

For $f \in \mathcal{A}$, the generalized differential operator $\mathcal{J}_{\gamma,\mu}^{m,\delta}: \mathcal{A}_\alpha \rightarrow \mathcal{A}_\alpha$ is defined

$$\begin{aligned}
\mathcal{J}_{\gamma,\mu}^{m,\delta} f(z)^\alpha &= \mathcal{J}(\mathcal{J}_{\gamma,\mu}^{m-1,\delta} f(z)^\alpha) \left(\frac{1-(\gamma-\delta)}{\mu+\delta} \right) (\mathcal{J}_{\gamma,\mu}^{m-1,\delta} f(z)^\alpha) + \frac{\gamma-\delta}{\mu+\delta} z (\mathcal{J}_{\gamma,\mu}^{m-1,\delta} f(z)^\alpha)' \quad (1.3) \\
&= \left(\frac{1+(1-\alpha)(\delta-\gamma)}{\mu+\delta} \right)^m z^\alpha + \sum_{k=2}^{\infty} \left(\frac{1+(1-\alpha)(\alpha+k-1)}{\mu+\delta} \right)^m a_k(\alpha) z^{\alpha+k-1},
\end{aligned}$$

and

where $(\mu > 0, \delta, \gamma \geq 0; m \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}; \alpha > 0, z \in \mathcal{U})$. For more details see [6].

From equation (1.3), we get

$$\begin{aligned}
\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z) &= \left(\frac{1-(\gamma-\delta)}{\mu+\delta} \right) (\mathcal{J}_{\gamma,\mu}^{m,\delta} f(z)) + \frac{\gamma-\delta}{\mu+\delta} (\mathcal{J}_{\gamma,\mu}^{m,\delta} f(z))' \\
\frac{\gamma-\delta}{\mu+\delta} z (\mathcal{J}_{\gamma,\mu}^{m,\delta} f(z))' &= \mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z) - \left(\frac{1-(\gamma-\delta)}{\mu+\delta} \right) (\mathcal{J}_{\gamma,\mu}^{m,\delta} f(z)). \quad (1.4)
\end{aligned}$$

We indicate by \mathcal{H} , the class of all function ϑ which are holomorphic and univalent in \mathcal{U} each of them $\vartheta(\mathcal{U})$ is convex such that $\vartheta(0) = 1$ and $\operatorname{Re}(\vartheta(z)) > 0, z \in \mathcal{U}$.

In this paper, we present differential operator from in relation to (1.1) and (1.2). We must use the principle of subordination between holomorphic functions to present the subclasses of starlike, convex and close-to-convex functions $S^*(\eta, \vartheta)$, $\mathcal{C}(\eta, \vartheta)$ and $K(\eta, \xi; \varphi, \vartheta)$ of order η respectively, for the function $\vartheta, \varphi \in \mathcal{H}$ which are defined by:

$$\begin{aligned}
S_{\alpha,\gamma,\mu}^{m,\delta}(\eta, \vartheta) &= \{f(z) \in \mathcal{A}: \mathcal{J}_{\gamma,\mu}^{m,\delta} f(z) \in S^*(\eta, \vartheta)\} \\
\mathcal{C}_{\alpha,\gamma,\mu}^{m,\delta}(\eta, \vartheta) &= \{f(z) \in \mathcal{A}: \mathcal{J}_{\gamma,\mu}^{m,\delta} f(z) \in \mathcal{C}(\eta, \vartheta)\}, \\
K_{\alpha,\gamma,\mu}^{m,\delta}(\eta, \xi; \varphi, \vartheta) &= \{f(z) \in \mathcal{A}: \mathcal{J}_{\gamma,\mu}^{m,\delta} f(z) \in K(\eta, \xi; \varphi, \vartheta)\}.
\end{aligned}$$

Next ,we will give introductory results that will be used in the show that of the main results of the paper.

Lemma:1.1.[1,2,10] Let ϖ be convex, univalent in \mathcal{U} with $\varpi(0) = 1$ and $\operatorname{Re}\{\pi\varpi(z) + \eta\} \geq 0, \pi, \eta \in \mathbb{C}$. If \mathcal{P} is holomorphic in \mathcal{U} with $\mathcal{P}(0) = 1$, then

$$\mathcal{P}(z) + \frac{z\mathcal{P}'(z)}{\pi\mathcal{P}(z)+\eta} < \varpi(z), z \in \mathcal{U}, \text{ means } \mathcal{P}(z) < \varpi(z), z \in \mathcal{U}.$$

Lemma:1.2.[8,10] Let ϖ be convex, univalent in \mathcal{U} and w be holomorphic in \mathcal{U} with $\operatorname{Re}(w(z)) \geq 0$. If \mathcal{P} is holomorphic in \mathcal{U} with $\mathcal{P}(0) = \varpi(0)$, then

$$\mathcal{P}(z) + w(z)z\mathcal{P}'(z) < \varpi(z), z \in \mathcal{U}, \text{ means } \mathcal{P}(z) < \varpi(z), z \in \mathcal{U}.$$

We will introduce some inclusion properties of the operator $\mathcal{J}_{\gamma,\mu}^{m,\delta} f(z)^\alpha$, by the *principle of subordination*.

2.Main Results

Theorem:2.1 Let $f \in \mathcal{A}$ and let $\vartheta \in \mathcal{H}$ with $\operatorname{Re}\left((1-\eta)\vartheta(z) + \eta + \frac{1-(\gamma-\delta)}{\mu+\delta}\right) > 0$. Then,

$$S_{\alpha,\gamma,\mu}^{m+1,\delta}(\eta, \vartheta) \subset S_{\alpha,\gamma,\mu}^{m,\delta}(\eta, \vartheta)$$

Proof: Let $f(z) \in S_{\alpha,\gamma,\mu}^{m+1,\delta}(z)$ and let

$$\mathcal{P}(z) = \frac{1}{1-\eta} \left(\frac{z(\mathcal{J}_{\gamma,\mu}^{m,\delta} f(z))'}{\mathcal{J}_{\gamma,\mu}^{m,\delta} f(z)} - \eta \right). \quad (2.1)$$

Applying (1.5) in (2.1), we get:

$$\begin{aligned} \mathcal{P}(z) &= \frac{1}{1-\eta} \left(\frac{\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z) - \left(\frac{1-(\gamma-\delta)}{\mu+\delta}\right) (\mathcal{J}_{\gamma,\mu}^{m,\delta} f(z))}{\frac{\lambda-\delta}{\mu+\delta} \mathcal{J}_{\gamma,\mu}^{m,\delta} f(z)} - \eta \right) \\ &= \left(\frac{\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z) - \left(\frac{1-(\gamma-\delta)}{\mu+\delta}\right) (\mathcal{J}_{\gamma,\mu}^{m,\delta} f(z))}{\frac{\lambda-\delta}{\mu+\delta} \mathcal{J}_{\gamma,\mu}^{m,\delta} f(z)} - \eta \right) = (1-\eta)\mathcal{P}(z) \end{aligned}$$

$$\left(\frac{\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z) - \left(\frac{1-(\gamma-\delta)}{\mu+\delta} \right) (\mathcal{J}_{\gamma,\mu}^{m,\delta} f(z))}{\frac{\gamma-\delta}{\mu+\delta} \mathcal{J}_{\gamma,\mu}^{m,\delta} f(z)} \right) = (1-\eta)\mathcal{P}(z) + \eta$$

$$\left(\frac{\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z)}{\frac{\gamma-\delta}{\mu+\delta} \mathcal{J}_{\gamma,\mu}^{m,\delta} f(z)} - \frac{\left(\frac{1-(\gamma-\delta)}{\mu+\delta} \right) (\mathcal{J}_{\gamma,\mu}^{m,\delta} f(z))}{\frac{\gamma-\delta}{\mu+\delta} \mathcal{J}_{\gamma,\mu}^{m,\delta} f(z)} \right) = (1-\eta)\mathcal{P}(z) + \eta$$

$$\frac{\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z)}{\frac{\gamma-\delta}{\mu+\delta} \mathcal{J}_{\gamma,\mu}^{m,\delta} f(z)} = (1-\eta)\mathcal{P}(z) + \eta + \left(\frac{1-(\gamma-\delta)}{\gamma-\delta} \right). \quad (2.2)$$

Differentiating (2.2) logarithmically with respect to z

$$\text{Log} \left(\frac{\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z)}{\frac{\gamma-\delta}{\mu+\delta} \mathcal{J}_{\gamma,\mu}^{m,\delta} f(z)} \right) = \text{Log} \left((1-\eta)\mathcal{P}(z) + \eta + \frac{1-(\gamma-\delta)}{\gamma-\delta} \right)$$

$$\text{Log} \left(\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z) \right) - \text{Log} \left(\frac{\gamma-\delta}{\mu+\delta} \mathcal{J}_{\gamma,\mu}^{m,\delta} f(z) \right) = \text{Log} \left((1-\eta)\mathcal{P}(z) + \eta + \frac{1-(\gamma-\delta)}{\gamma-\delta} \right)$$

$$\frac{(\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z))'}{(\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z))} - \frac{\left(\frac{\gamma-\delta}{\mu+\delta} \right) (\mathcal{J}_{\gamma,\mu}^{m,\delta} f(z))'}{\left(\frac{\gamma-\delta}{\mu+\delta} \right) (\mathcal{J}_{\gamma,\mu}^{m,\delta} f(z))} = \frac{(1-\eta)\mathcal{P}'(z)}{(1-\eta)\mathcal{P}(z) + \frac{1-(\gamma-\delta)}{\gamma-\delta}}$$

$$\frac{(\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z))'}{(\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z))} = \frac{(\mathcal{J}_{\gamma,\mu}^{m,\delta} f(z))'}{(\mathcal{J}_{\gamma,\mu}^{m,\delta} f(z))} + \frac{(1-\eta)\mathcal{P}'(z)}{(1-\eta)\mathcal{P}(z) + \frac{1-(\gamma-\delta)}{\gamma-\delta}}. \quad (2.3)$$

On the other hand:

$$\frac{(\mathcal{J}_{\gamma,\mu}^{m,\delta} f(z))'}{(\mathcal{J}_{\gamma,\mu}^{m,\delta} f(z))} = \frac{(1-\eta)\mathcal{P}(z) + \eta}{z}, \quad (2.4)$$

using (2.3) and (2.4)

$$\frac{(\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z))'}{(\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z))} = \frac{(1-\eta)\mathcal{P}(z) + \eta}{z} + \frac{(1-\eta)\mathcal{P}'(z)}{(1-\eta)\mathcal{P}(z) + \frac{1-(\gamma-\delta)}{\gamma-\delta}}$$

$$\frac{(\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z))'}{(\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z))} = \frac{(1-\eta)\mathcal{P}(z)}{z} + \frac{\eta}{z} + \frac{(1-\eta)\mathcal{P}'(z)}{(1-\eta)\mathcal{P}(z) + \frac{1-(\gamma-\delta)}{\gamma-\delta}}$$

$$\begin{aligned} \frac{(\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z))'}{(\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z))} - \frac{\eta}{z} &= \frac{(1-\eta)\mathcal{P}(z)}{z} + \frac{(1-\eta)\mathcal{P}'(z)}{(1-\eta)\mathcal{P}(z) + \frac{1-(\gamma-\delta)}{\gamma-\delta}} \\ \frac{z(\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z))'}{(\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z))} - \eta &= \frac{(1-\eta)\left[\mathcal{P}(z)\left((1-\eta)\mathcal{P}(z) + \eta + \frac{1-(\gamma-\delta)}{\gamma-\delta}\right)\right] + z\mathcal{P}'(z)}{\left[(1-\eta)\mathcal{P}(z) + \eta + \frac{1-(\gamma-\delta)}{\gamma-\delta}\right]} \\ \frac{1}{1-\eta} \left(\frac{z(\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z))'}{(\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z))} - \eta \right) &= \frac{\mathcal{P}(z)\left((1-\eta)\mathcal{P}(z) + \eta + \frac{1-(\gamma-\delta)}{\gamma-\delta}\right)}{\left((1-\eta)\mathcal{P}(z) + \eta + \frac{1-(\gamma-\delta)}{\gamma-\delta}\right)} + \frac{z\mathcal{P}'(z)}{\left((1-\eta)\mathcal{P}(z) + \eta + \frac{1-(\gamma-\delta)}{\gamma-\delta}\right)} \\ \frac{1}{1-\eta} \left(\frac{z(\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z))'}{(\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z))} - \eta \right) &= \mathcal{P}(z) + \frac{z\mathcal{P}'(z)}{\left((1-\eta)\mathcal{P}(z) + \eta + \frac{1-(\gamma-\delta)}{\gamma-\delta}\right)}. \end{aligned} \tag{2.5}$$

Applying Lemma (1.1) to (2.5),we get

$$\operatorname{Re}\{\pi\vartheta(z) + \tau\} \geq 0,$$

$$\pi\operatorname{Re}(\vartheta(z)) + \tau \geq 0$$

Since ϑ convex, order τ

$$\operatorname{Re}\left\{1 + \frac{zf''(z)}{f'(z)}\right\} > \pi \left| \frac{zf''(z)}{f'(z)} \right| + \tau, z \in \mathcal{U}, 0 \leq \tau < 1, \pi \geq 0, \pi + \tau \geq 0$$

$$\operatorname{Re}\left\{(1-\eta)\vartheta(z) + \eta + \frac{1-(\gamma-\delta)}{\gamma-\delta}\right\} > 0, \delta, \gamma \geq 0$$

$$(1-\eta)\operatorname{Re}(\vartheta(z)) + \eta + \frac{1-(\gamma-\delta)}{\gamma-\delta} > 0, \vartheta(0) = 1, \operatorname{Re}(\vartheta(z)) > 0, z \in \mathcal{U},$$

$$(1-\eta) + \eta + \frac{1-(\gamma-\delta)}{\gamma-\delta} > 0,$$

then

$$\mathcal{P}(z) + \frac{z\mathcal{P}'(z)}{(1-\eta)\mathcal{P}(z) + \eta + \frac{1-(\gamma-\delta)}{\gamma-\delta}} < \vartheta(z), z \in \mathcal{U}, \text{ means } \mathcal{P}(z) < \vartheta(z), z \in \mathcal{U},$$

i.e. $f(z) \in \mathcal{J}_{\gamma,\mu}^{m,\delta} f(z)$.

Thus,

$$S_{\alpha,\gamma,\mu}^{m+1,\delta} f(z) \subset S_{\alpha,\gamma,\mu}^{m,\delta} f(z),$$

□

Theorem:2.2. Let $f \in A$ and let $\vartheta \in \mathcal{H}$ with $\operatorname{Re}\left\{(1-\eta)\vartheta(z) + \eta + \frac{1-(\gamma-\delta)}{\gamma-\delta}\right\} > 0$. Then,

$$C_{\alpha,\gamma,\mu}^{m+1,\delta} f(z) \subset C_{\alpha,\gamma,\mu}^{m,\delta} f(z).$$

Proof: From Remark 1.1, we have: $f \in \mathcal{A}$ is starlike and convex, respectively, order τ , if and only if

$$\operatorname{Re} \left\{ \frac{zf'(z)}{f(z)} \right\} > \tau, 0 \leq \tau < 1, |z| < 1,$$

and

$$\operatorname{Re} \left\{ 1 + \frac{zf''(z)}{f'(z)} \right\} > \tau, 0 \leq \tau < 1, |z| < 1.$$

A function $f \in \mathcal{A}$ is uniformly starlike and uniformly convex, respectively, of order τ , if and only if [5]

$$\operatorname{Re} \left\{ \frac{zf'(z)}{f(z)} \right\} > \pi \left| \frac{zf'(z)}{f(z)} - 1 \right| + \tau, z \in \mathcal{U}$$

and

$$\operatorname{Re} \left\{ 1 + \frac{zf''(z)}{f'(z)} \right\} > \pi \left| \frac{zf''(z)}{f'(z)} \right| + \tau, z \in \mathcal{U},$$

where $\pi \geq 0$, $0 \leq \tau < 1$ and $\pi + \tau \geq 0$.

The relation between the classes of starlike and convex functions, obviously, led us to the next relation .

$$f \in C_{\alpha,\gamma,\mu}^{m+1,\delta}(\eta, \vartheta) \Leftrightarrow zf' \in S_{\alpha,\gamma,\mu}^{m+1,\delta}(\eta, \vartheta).$$

From Theorem 2.1, we get

$$\begin{aligned} f \in C_{\alpha,\gamma,\mu}^{m+1,\delta}(\eta, \vartheta) &\Leftrightarrow zf' \in S_{\alpha,\gamma,\mu}^{m+1,\delta}(\eta, \vartheta) \subset S_{\alpha,\gamma,\mu}^{m,\delta}(\eta, \vartheta) \\ &\Rightarrow zf' \in S_{\alpha,\gamma,\mu}^{m,\delta}(\eta, \vartheta) \\ &\Rightarrow zf' \in C_{\alpha,\gamma,\mu}^{m,\delta}(\eta, \vartheta). \end{aligned}$$

Thus,

$$C_{\alpha,\gamma,\mu}^{m+1,\delta}(\eta, \vartheta) \subset C_{\alpha,\gamma,\mu}^{m,\delta}(\eta, \vartheta).$$

□

The function $\vartheta(z) = \frac{1-Az}{1-Bz}$ is holomorphic and satisfies $\vartheta(0) = 1$. Thus, we get the next corollaries:

Corollary:2.1. Let $f \in \mathcal{A}$ and $\vartheta = \frac{1+Az}{1-Bz}$, $-1 \leq B \leq A \leq 1$ in Theorem 2.1. Then:

$$S_{\alpha,\gamma,\mu}^{m+1,\delta}(\eta, \mathcal{A}, \mathcal{B}) \subset S_{\alpha,\gamma,\mu}^{m,\delta}(\eta, \mathcal{A}, \mathcal{B}).$$

Corollary:2.2. Let $f \in \mathcal{A}$ and $\vartheta = \frac{1+Az}{1-Bz}$, $-1 \leq B \leq A \leq 1$ in Theorem 2.1. Then:

$$K_{\alpha,\gamma,\mu}^{m+1,\delta}(\eta, \mathcal{A}, \mathcal{B}) \subset K_{\alpha,\gamma,\mu}^{m,\delta}(\eta, \mathcal{A}, \mathcal{B}).$$

Theorem:2.3 Let $f \in \mathcal{A}$ and let $\vartheta, \varphi \in \mathcal{H}$ with $\operatorname{Re} \left\{ (1 - \eta)\vartheta(z) + \eta + \frac{1-(\gamma-\delta)}{\gamma-\delta} \right\} > 0$. Then:

$$K_{\alpha, \gamma, \mu}^{m+1, \delta}(\eta, \xi; \vartheta, \varphi) \subset K_{\alpha, \gamma, \mu}^{m, \delta}(\eta, \xi; \vartheta, \varphi).$$

Proof : Let $f \in K_{\alpha, \gamma, \mu}^{m+1, \delta}(\eta, \xi; \vartheta, \varphi)$, and let $g \in S_{\alpha, \gamma, \mu}^{m+1, \delta}(\eta, \vartheta)$ such that:

$$\operatorname{Re} \left\{ \frac{z \left(\mathcal{J}_{\gamma, \mu}^{m+1, \delta} f(z) \right)'}{\mathcal{J}_{\gamma, \mu}^{m+1, \delta} g(z)} \right\} > \xi, z \in U.$$

The next should, we get

$$\frac{1}{1 - \xi} \left(\frac{z \left(\mathcal{J}_{\gamma, \mu}^{m+1, \delta} f(z) \right)'}{\mathcal{J}_{\gamma, \mu}^{m+1, \delta} g(z)} - \xi \right) < \varphi, z \in \mathcal{U}.$$

Let

$$\mathcal{P}(z) = \frac{1}{1 - \xi} \left(\frac{z \left(\mathcal{J}_{\gamma, \mu}^{m+1, \delta} f(z) \right)'}{\mathcal{J}_{\gamma, \mu}^{m+1, \delta} g(z)} - \xi \right) \quad (2.6)$$

$$\left(\frac{z \left(\mathcal{J}_{\gamma, \mu}^{m+1, \delta} f(z) \right)'}{\mathcal{J}_{\gamma, \mu}^{m+1, \delta} g(z)} - \xi \right) = (1 - \xi) \mathcal{P}(z)$$

$$\frac{z \left(\mathcal{J}_{\gamma, \mu}^{m+1, \delta} f(z) \right)'}{\mathcal{J}_{\gamma, \mu}^{m+1, \delta} g(z)} = (1 - \xi) \mathcal{P}(z) + \xi.$$

From (1.6), we get

$$\frac{\mathcal{J}_{\gamma, \mu}^{m+1, \delta} f(z) - \left(\frac{1-(\gamma-\delta)}{\mu-\delta} \right) \left(\mathcal{J}_{\gamma, \mu}^{m, \delta} f(z) \right)}{\frac{\gamma-\delta}{\mu+\delta} \mathcal{J}_{\gamma, \mu}^{m, \delta} g(z)} = (1 - \xi) \mathcal{P}(z) + \xi$$

$$\frac{\mathcal{J}_{\gamma, \mu}^{m+1, \delta} f(z) - \left(\frac{1-(\gamma-\delta)}{\mu-\delta} \right) \left(\mathcal{J}_{\gamma, \mu}^{m, \delta} f(z) \right)}{\frac{\gamma-\delta}{\mu+\delta}} = [(1 - \xi) \mathcal{P}(z) + \xi] \left(\mathcal{J}_{\gamma, \mu}^{m, \delta} g(z) \right)$$

$$\frac{\mathcal{J}_{\gamma, \mu}^{m+1, \delta} f(z)}{\frac{\gamma-\delta}{\mu+\delta}} - \frac{\left(\frac{1-(\gamma-\delta)}{\mu-\delta} \right) \left(\mathcal{J}_{\gamma, \mu}^{m, \delta} f(z) \right)}{\frac{\gamma-\delta}{\mu+\delta}} = [(1 - \xi) \mathcal{P}(z) + \xi] \left(\mathcal{J}_{\gamma, \mu}^{m, \delta} g(z) \right).$$

Mean that

$$\begin{aligned} \frac{(\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z))'}{\frac{\gamma-\delta}{\mu+\delta}} &= \left(\frac{1-(\gamma-\delta)}{\mu-\delta} \right) (\mathcal{J}_{\gamma,\mu}^{m,\delta} f(z))' + [(1-\xi)\mathcal{P}(z) + \xi](\mathcal{J}_{\gamma,\mu}^{m,\delta} g(z))' \\ &\quad + [(1-\xi)\mathcal{P}'(z)](\mathcal{J}_{\gamma,\mu}^{m,\delta} g(z)). \\ \frac{z(\mu+\delta)}{\gamma-\delta} (\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z))' &= \left(\frac{1-(\gamma-\delta)}{\gamma-\delta} \right) z(\mathcal{J}_{\gamma,\mu}^{m,\delta} f(z))' + [(1-\xi)\mathcal{P}(z) + \xi]z(\mathcal{J}_{\gamma,\mu}^{m,\delta} g(z))' \\ &\quad + [(1-\xi)z\mathcal{P}'(z)](\mathcal{J}_{\gamma,\mu}^{m,\delta} g(z)). \end{aligned} \quad (2.7)$$

Also, by Theorem 2.1, $g \in S_{\gamma,\mu}^{m+1,\delta}(\eta, \vartheta) \rightarrow g \in S_{\gamma,\mu}^{m,\delta}(\eta, \vartheta)$.

Thus, let

$$q(z) = \frac{1}{1-\eta} \left(\frac{z(\mathcal{J}_{\gamma,\mu}^{m,\delta} g(z))'}{\mathcal{J}_{\gamma,\mu}^{m,\delta} g(z)} - \eta \right). \quad (2.8)$$

Using (1.6) in (2.8), we get

$$\begin{aligned} \left(\frac{z(\mathcal{J}_{\gamma,\mu}^{m,\delta} g(z))'}{\mathcal{J}_{\gamma,\mu}^{m,\delta} g(z)} \right)' &= (1-\eta)q(z) + \eta \\ \frac{\mathcal{J}_{\gamma,\mu}^{m+1,\delta} g(z) - \left(\frac{1-(\gamma-\delta)}{\mu-\delta} \right) (\mathcal{J}_{\gamma,\mu}^{m,\delta} g(z))}{\frac{\gamma-\delta}{\mu+\delta} \mathcal{J}_{\gamma,\mu}^{m,\delta} g(z)} &= (1-\eta)q(z) + \eta \\ \frac{\mathcal{J}_{\gamma,\mu}^{m+1,\delta} g(z)}{\frac{\gamma-\delta}{\mu+\delta} \mathcal{J}_{\gamma,\mu}^{m,\delta} g(z)} - \frac{\left(\frac{1-(\gamma-\delta)}{\mu-\delta} \right) (\mathcal{J}_{\gamma,\mu}^{m,\delta} g(z))}{\frac{\gamma-\delta}{\mu+\delta} \mathcal{J}_{\gamma,\mu}^{m,\delta} g(z)} &= (1-\eta)q(z) + \eta \\ \frac{\mathcal{J}_{\gamma,\mu}^{m+1,\delta} g(z)}{\frac{\gamma-\delta}{\mu+\delta} \mathcal{J}_{\gamma,\mu}^{m,\delta} g(z)} &= (1-\eta)q(z) + \eta + \frac{(1-(\gamma-\delta))}{\gamma-\delta}. \end{aligned} \quad (2.9)$$

And further, from (2.7) and (2.9), we get

$$\begin{aligned} \frac{\frac{\mu+\delta}{\gamma-\delta} (\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z))'}{\frac{\mu+\delta}{\gamma-\delta} \frac{\mathcal{J}_{\gamma,\mu}^{m+1,\delta} g(z)}{\mathcal{J}_{\gamma,\mu}^{m,\delta} g(z)}} &= \frac{\left(\frac{1-(\gamma-\delta)}{\gamma-\delta} \right) z(\mathcal{J}_{\gamma,\mu}^{m,\delta} f(z))' + [(1-\xi)\mathcal{P}(z) + \xi]z(\mathcal{J}_{\gamma,\mu}^{m,\delta} g(z))'}{\left[(1-\eta)q(z) + \eta + \frac{(1-(\gamma-\delta))}{\gamma-\delta} \right]} \\ &\quad + \frac{[(1-\xi)z\mathcal{P}'(z)](\mathcal{J}_{\gamma,\mu}^{m,\delta} g(z))}{\left[(1-\eta)q(z) + \eta + \frac{(1-(\gamma-\delta))}{\gamma-\delta} \right]} \end{aligned}$$

$$\begin{aligned}
 \frac{\frac{\mu+\delta}{\gamma-\delta} z(\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z))'}{\frac{\mu+\delta}{\gamma-\delta} (\mathcal{J}_{\gamma,\mu}^{m+1,\delta} g(z))} &= \frac{\left(\frac{1-(\gamma-\delta)}{\gamma-\delta}\right) z(\mathcal{J}_{\gamma,\mu}^{m,\delta} f(z))' + [(1-\xi)\mathcal{P}(z) + \xi]z(\mathcal{J}_{\gamma,\mu}^{m,\delta} g(z))'}{(\mathcal{J}_{\gamma,\mu}^{m,\delta} g(z)) \left[(1-\eta)q(z) + \eta + \frac{(1-(\gamma-\delta))}{\gamma-\delta}\right]} \\
 &+ \frac{[(1-\xi)z\mathcal{P}'(z)](\mathcal{J}_{\gamma,\mu}^{m,\delta} g(z))}{(\mathcal{J}_{\gamma,\mu}^{m,\delta} g(z)) \left[(1-\eta)q(z) + \eta + \frac{(1-(\gamma-\delta))}{\gamma-\delta}\right]} \\
 &= \frac{\left(\frac{1-(\gamma-\delta)}{\gamma-\delta}\right) z(\mathcal{J}_{\gamma,\mu}^{m,\delta} f(z))'}{(\mathcal{J}_{\gamma,\mu}^{m,\delta} g(z)) \left[(1-\eta)q(z) + \eta + \frac{(1-(\gamma-\delta))}{\gamma-\delta}\right]} + \frac{[(1-\xi)\mathcal{P}(z) + \xi]z\left((\mathcal{J}_{\gamma,\mu}^{m,\delta} g(z))'\right)}{(\mathcal{J}_{\gamma,\mu}^{m,\delta} g(z)) \left[(1-\eta)q(z) + \eta + \frac{(1-(\gamma-\delta))}{\gamma-\delta}\right]} \\
 &+ \frac{[(1-\xi)z\mathcal{P}'(z)](\mathcal{J}_{\gamma,\mu}^{m,\delta} g(z))}{(\mathcal{J}_{\gamma,\mu}^{m,\delta} g(z)) \left[(1-\eta)q(z) + \eta + \frac{(1-(\gamma-\delta))}{\gamma-\delta}\right]} \\
 &= \frac{\frac{(1-(\gamma-\delta))}{\gamma-\delta} [(1-\xi)\mathcal{P}(z) + \xi]}{\left[(1-\eta)q(z) + \eta + \frac{(1-(\gamma-\delta))}{\gamma-\delta}\right]} + \frac{[(1-\xi)\mathcal{P}(z) + \xi][(1-\eta)q(z) + \eta]}{\left[(1-\eta)q(z) + \eta + \frac{(1-(\gamma-\delta))}{\gamma-\delta}\right]} + \frac{[(1-\xi)z\mathcal{P}'(z)]}{\left[(1-\eta)q(z) + \eta + \frac{(1-(\gamma-\delta))}{\gamma-\delta}\right]} \\
 &= \frac{\left(\frac{1-(\gamma-\delta)}{\gamma-\delta}\right) [(1-\xi)\mathcal{P}(z) + \xi] + [(1-\xi)\mathcal{P}(z) + \xi][(1-\eta)q(z) + \eta]}{\left[(1-\eta)q(z) + \eta + \frac{(1-(\gamma-\delta))}{\gamma-\delta}\right]} + \frac{[(1-\xi)z\mathcal{P}'(z)]}{\left[(1-\eta)q(z) + \eta + \frac{(1-(\gamma-\delta))}{\gamma-\delta}\right]} \\
 \frac{z(\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z))'}{(\mathcal{J}_{\gamma,\mu}^{m+1,\delta} g(z))} &= [(1-\xi)\mathcal{P}(z) + \xi] + \frac{[(1-\xi)z\mathcal{P}'(z)]}{\left[(1-\eta)q(z) + \eta + \frac{(1-(\gamma-\delta))}{\gamma-\delta}\right]}. \tag{2.10}
 \end{aligned}$$

Algebraic manipulation in (2.10) gives:

$$\begin{aligned}
 \frac{z(\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z))'}{(\mathcal{J}_{\gamma,\mu}^{m+1,\delta} g(z))} - \xi &= [(1-\xi)\mathcal{P}(z)] + \frac{[(1-\xi)z\mathcal{P}'(z)]}{\left[(1-\eta)q(z) + \eta + \frac{(1-(\gamma-\delta))}{\gamma-\delta}\right]} \\
 \frac{1}{1-\xi} \left(\frac{z(\mathcal{J}_{\gamma,\mu}^{m+1,\delta} f(z))'}{(\mathcal{J}_{\gamma,\mu}^{m+1,\delta} g(z))} - \xi \right) &= \mathcal{P}(z) + \frac{[z\mathcal{P}'(z)]}{\left[(1-\eta)q(z) + \eta + \frac{(1-(\gamma-\delta))}{\gamma-\delta}\right]}.
 \end{aligned}$$

So, we get

$$w(z) = \frac{1}{\left[(1 - \eta)q(z) + \eta + \frac{(1-(\gamma-\delta))}{\gamma-\delta} \right]}$$

Applying Lemma 1.2, we get

$$\mathcal{P}(z) < \varphi(z), \text{ mean that } f \in K_{\gamma, \mu}^{m+1, \delta}(\eta, \xi; \vartheta, \varphi) \quad \square$$

3. Conclusion: In this article we concluded that in this case of applying the differential operator for univalent function using a Faber polynomial it remains preserving it's geometric properties and there are other conditions added to obtain results inside the unit disk.

REFERENCES

- [1] Bulboaca, T. (2005). Differential Subordinations and Superordinations: Recent results. Casa Cartii de Stiinta
- [2] Choi, J. H., Saigo, M., & Srivastava, H. M. (2002). Some inclusion properties of a certain family of integral operators. Journal of Mathematical Analysis and Applications, 276(1), 432-445.
- [3] Duren, P. L. (2001). Univalent functions (Vol. 259). Springer Science & Business Media.
- [4] Eenigenburg, P., Mocanu, P. T., Miller, S. S., & Reade, M. O. (1983). On a Briot-Bouquet differential subordination. In General Inequalities 3 (pp. 339-348). Birkhauser, Basel.
- [5] Goodman, A. W. (1991). On uniformly convex functions. In Annales Polonici Mathematici (Vol. 56, No. 1, pp. 87-92).
- [6] Juma, A. R. S., Abdulhussain, M. S., & AL-khafaji, S. N. (2019). Faber Polynomial Coefficient Estimates for Subclass of Analytic Bi-Bazilevic Functions Defined by Differential Operator. Baghdad Science Journal, 16(1 Supplement), 248-253.
- [7] Makinde, D. O. (2018). A new multiplier differential operator. Adv. Math., Sci. J, 7, 109-114.
- [8] Miller, S. S. Mocanu, P. T. (1981). Differential Subordinations and univalent functions. The Michigan Mathematical Journal, 28(2), 157-172.
- [9] Shigeyoshi, O. (Ed.). (1992). Current topics in analytic theory. world Scientific.
- [10] Swamy, S. R. (2012). Inclusion properties of certain subclasses of analytic functions. Int. Math. Forum (Vol. 7, No. 36, pp. 1751-1760).