

1-7-2020

## Roles of Superoxide dismutase (SOD), Malondialdehyde (MDA), 8-iso-prostaglandin F<sub>2</sub>α ( 8-iso-PGF<sub>2</sub>α) as oxidative stress in development and progression of Breast cancer in Iraqi females patients

Ferdous Abbas Jabir

*Department of biochemistry, College of medicine, Univesity of Al-Qadisiyah, Iraq*

Ahmed Sabah Shaker

*Department of biochemistry, College of medicine, Univesity of Al-Qadisiyah, Iraq,*

*ahmedalyssain@gmail.com*

Follow this and additional works at: <https://qjps.researchcommons.org/home>

 Part of the [Chemistry Commons](#)

---

### Recommended Citation

Jabir, Ferdous Abbas and Shaker, Ahmed Sabah (2020) "Roles of Superoxide dismutase (SOD), Malondialdehyde (MDA), 8-iso-prostaglandin F<sub>2</sub>α ( 8-iso-PGF<sub>2</sub>α) as oxidative stress in development and progression of Breast cancer in Iraqi females patients," *Al-Qadisiyah Journal of Pure Science*: Vol. 25: No. 1, Article 13.

DOI: 10.29350/2411-3514.1218

Available at: <https://qjps.researchcommons.org/home/vol25/iss1/13>

This Article is brought to you for free and open access by Al-Qadisiyah Journal of Pure Science. It has been accepted for inclusion in Al-Qadisiyah Journal of Pure Science by an authorized editor of Al-Qadisiyah Journal of Pure Science. For more information, please contact [bassam.alfarhani@qu.edu.iq](mailto:bassam.alfarhani@qu.edu.iq).

Received : 16/12/2019

Accepted :13/1/2020

## Roles of Superoxide dismutase(SOD) , Malondialdehyde (MDA), 8-iso-prostaglandinF2 $\alpha$ ( 8-iso-PGF2 $\alpha$ )as oxidative stress in development and progression of Brest cancer in Iraqi females patients

Ferdous Abbas Jabir<sup>1</sup>, Ahmed Sabah Shaker<sup>2\*</sup><sup>1</sup> Department of bio chemistry, College of medicine, Univesity of Al-Qadisiyah, Iraq<sup>2\*</sup>corresponding author, email: ahmedalyssain@gmail.com**Abstract:-**

Oxidative stress occurs as a result of disturbance in the balance between the production of reactive oxygen species (free radicals) and antioxidant defenses. Markers of oxidative stress were measured the markers of oxidative stress in breast cancer patients after diagnosis of breast cancer and compared these plasma blood levels controls This study was conducted to three markers of oxidative stress ;these are (SOD) enzyme ,malondialdehyde (MDA)and8-iso-prostaglandinF2 $\alpha$  plasma of patients with breast cancer and compare with controls .In this study ; the mean MDA (ng/ml) levels for the breast cancer patients and the controls were55.91 $\pm$ 3.31 and40.61 $\pm$ 3.76 respectively, while the SOD (pg/ml) levels were1530.37 $\pm$ 80.4 and1851.4 9 $\pm$ 93.65 respectively and the 8-iso-PGF2 $\alpha$  (ng/ml ) levels were 40.16 $\pm$ 3.31 and 30.16 $\pm$ 2.34 difference of the mean were statistically significant (p value <0.05).

**Keywords:** Breast cancer, Malondialdehyde, 8-iso-PGF2 $\alpha$ , Oxidative stress, Superoxide dismutase

**Introduction:-**

Breast cancer is that the most typically diagnosed cancer among women within the overwhelming majority (140/184) of states worldwide, representing 1 / 4 of all cancers diagnosed in girls(DeSantis et al., 2015). Breast cancer is the formation of a malignancy that has developed from cells in the breast(Israyelyan, 2003). Breast most cancer is a multifactorial disease environmental elements certain as radiation, viruses or chemical compounds bear a vital function so carcinogenic agents. At the identical time, numerous secondary factors: gender, age, economic development regarding the country regarding residence,(Tillashaykhov, Khudaykulov, Danilova, Usupbekov, & Alimkhodjaeva, 2017). Oxidative stress defined as an imbalance of oxidants and antioxidants in favor on the oxidants. the antioxidant defense mechanisms on the ethnic body, oxidative damage perform happen in imitation of lipids, proteins, and immediately to

DNA (Lee, Cai, Shu, & Nechuta, 2017). It has been hypothesized as operative oxygen species (ROS) oxidative power may also play a critical role into cancer root cause and a progression or be associated along the proliferation potencies of breast most cancers (Hasan, Mathkor, & Al-Habal, 2012). Reactive oxygen species are produced consistently during normal metabolism, but are neutralized by antioxidants. However, when there is additional production regarding ROS and deficiency in an antioxidant system, oxidative stress occurs. raised DNA damage and mutations in genes (Kang, 2002). Free Radicals (oxidants ) In non-bonding electrons, free radicals tend to interact with other compounds and act as electronic receptors or oxidizing agents. Major oxidants include reactive oxygen species (ROS), nitrogen (RNS), RCS chloride), and sulfur. The most important antioxidant and the main cause of oxidative damage to the vital molecules in the body is ROS. ROS also plays an important role in the formation of interactive species such as

RNS(Nourazarian, Kangari, & Salmaninejad, 2014). One possible clinical application of oxidative stress in cancer is the use of oxidative stress markers as tumor markers. For example, it has been suggested enzyme concerning the nearly important enzymatic antioxidants (Hasan et al., 2012). There are twins main types about SOD in human cells: SOD (Mn-SOD), which is principally localized in PGF2 $\alpha$  (ng/ml ) were 40.16 $\pm$ 3.31 and 30.16 $\pm$ 2.34respectively . The difference of the mean between the MDA, SOD and 8-iso-PGF2 $\alpha$  levels of the breast cancer patients and the controls were be statistically significant (p value <0.05

**Table(1)oxidative stress in patients suffering from breast cancer**

Concentration	Patients	Controls	P value
SOD(pg/ml) (mean $\pm$ SD)	1530.37 $\pm$ 80.4	1851.4 9 $\pm$ 93.65	0.022
MDA(ng/ml) (mean $\pm$ SD)	55.91 $\pm$ 3.31	40.61 $\pm$ 3.76	0.008
PGF2a(ng/ml) (mean $\pm$ SD)	40.16 $\pm$ 3.31	30.16 $\pm$ 2.34	0.024

\* p<0.05 significantly raised in Breast Cancer

**Table(2) The mean age of the breast cancer patients and control**

Subjects	Age (mean $\pm$ SD)	P value
Control( n=30)	35 $\pm$ 3.94	0.360
patients( n=30)	40.45 $\pm$ 5.12	

(P>0.05) denote to non-significant .

SOD location key position within the anti oxidative network is of marked pathophysiological importance<sup>(Hasan et al., 2012)</sup>. It was found that this antioxidant phase decreased in breast cancer patients due to increased reactive oxygen species with comparative control (Sahu et al., 2015). These results have shown that very weak

that the blood SOD stage in breast cancer may be useful as a marker for the tumor (Noda & Wakasugi, 2001).Dismutase superoxide(SOD)

mitochondria, or Cu/Zncontaining SOD (Cu/Zn-SOD), which is chiefly localized in the cytosol. Mn-SOD (Er, Hou, Tsa, Lee, & Tsai, 2004).

Malondialdehyde(MDA)enzyme of the almost vital on these is (MDA), low-molecular-weight aldehydes derived beside lipid peroxidation processes, (Akbulut, Akbulut, Icli, & Büyükcelik, 2003). Malondialdehyde (MDA), the end-product over macromolecule peroxidation LPO) turning out beside the broad basic degradation over unsaturated swollen acids, do reason cross-linking among lipids, proteins, and nucleic acids (Pande, Negi, Khanna, Khanna, & Khanna, 2011). 8-iso-prostaglandinF2 $\alpha$  (8-iso-PGF2 $\alpha$ ) a gaggle of prostaglandin-like compounds, area unit biosynthesized chiefly from esterified arachidonic acid through a catalyst free radical–catalyzed reaction in vivo that has opened a unique facet of free-radical biology. (PGF2 $\alpha$ ), apotent prostaglandin that regulate several feminine hormones is related with breast cancer(Basu et al., 2016).

#### EXPERIMENTAL:-

Blood samples were collected from (30) patients with breast cancer in age (25-80) years and (30) healthy persons as a control group in age (25-80)years . five milliliters of venous blood was taken for each sample and collected in heparinized tube, then centrifuged at 3000 x g for 10 min (Sahu, Varma, & Kachhawa, 2015). Patients with other diseases such as blood pressure, diabetes and other diseases were excluded .In this study determination of plasma MDA ,SOD and8-iso-PGF2 $\alpha$  levels used MDA ,SOD and iso-PGF2 $\alpha$  kits by used Elisa Technique ( Elabscience. USA) . The data for biochemical analyses are expressed as mean and standard deviation (SD). Statistical comparisons were performed by Student's t-test using the SPSS version

#### Results and Dissection:-

The mean age of the breast cancer patients was 40.45 $\pm$ 5.12 years and those of the control group was 35 $\pm$ 3.94 years respectively. There was statistically non a significant. The MDA (ng/ml) levels for the breast cancer patients and the controls were 55.91 $\pm$ 3.31and 40.61 $\pm$ 3.76 respectively .while the SOD (pg/ml) levels were 1530.37 $\pm$ 80.4 and 1851.4 9 $\pm$ 93.65 and the 8-iso-

polyunsaturated fatty acids. a plus Plasma MDA levels have been reported in breast cancer. Our The results showed an increase in the level of MDA in breast cancer Compared with the controls thus agree with the former Studies and therefore suggest an increase in lipid peroxide In breast cancer patients(Gupta et al., 2012). The role of free radicals, oxidative stress, and lipid peroxidation in carcinogenesis and their contribution to the initiation and progression of the process are well documented . In recent years, using as8-iso-PGF2 $\alpha$  a marker of oxidative stress. results showed increase in as8-iso-PGF2 $\alpha$  level in breast cancer as compared to controls thus agreeing with the previous studies(Bennett et al., 1989; Dai & Zhu, 2009; Mannello et al., 2007),(Dai & Zhu, 2009) and thus suggesting increased lipid peroxidation in breast cancer patients(Gupta et al., 2012).

**4.Conclusion :**It found that there was an increase in the concentration of oxidative stress in patients with breast cancer. The MDA (ng/ml) statistically significant(0.008<0.05), the 8-iso-PGF2 $\alpha$  (ng/ml ) ( 0.024 <0.05) and the SOD ( 0.022<0.05) statistically significant .

#### References

- Elyasinia, F., Chegini, V., Olfat-Bakhsh, A., Pasalar, P., & Aminian, A. (2014). Superoxide dismutase activities in plasma of patients with breast cancer. *Archives of Breast Cancer, 1*(2), 69-72.
- Er, T.-K., Hou, M.-F., Tsa, E.-M., Lee, J.-N., & Tsai, L.-Y. (2004). Differential expression of manganese containing superoxide dismutase in patients with breast cancer in Taiwan. *Annals of Clinical & Laboratory Science, 34*(2), 159-164.
- Gönenç, A., Özkan, Y., Torun, M., & Şimşek, B. (2001). Plasma malondialdehyde (MDA) levels in breast and lung cancer patients. *Journal of clinical pharmacy and therapeutics, 26*(2), 141-144.
- Gupta, R. K., Patel, A. K., Kumari, R., Chugh, S., Shrivastav, C., Mehra, S., & Sharma, A. N. (2012). Interactions between oxidative stress, lipid profile and antioxidants in breast cancer: a case control study. *Asian Pacific Journal of Cancer Prevention, 13*(12), 6295-6298.
- Hasan, H. R., Mathkor, T. H., & Al-Habal, M. H. (2012). Superoxide dismutase isoenzyme activities in plasma and tissues of Iraqi patients with breast cancer. *Asian Pacific Journal of Cancer Prevention, 13*(6), 2571-2576.
- inhibitor defense systems in patients with malignant breast growth reduces the level of SOD enzymes cells to synthesize free radicals. Associate in nursing imbalance between the production of free radical enzyme and the defense mechanisms that have long pursued them is believed to be an important issue in nursing cause pathological process of cancer. In addition, we tend to find a statistically vital association between SOD and growth stage and grade (Elyasinia, Chegini, Olfat-Bakhsh, Pasalar, & Aminian, 2014). MDA, one of each lipid peroxide decomposition product, is believed in human plasma and has biological properties, is involved in many disease processes as well as cancer. Several studies have examined the opportunity related to the relationship between lipid peroxidation after cancer and advanced MDA levels in patients with cancer which have been presented in addition to evidence of this relationship (Gönenç, Özkan, Torun, & Şimşek, 2001). The role of free radicals, oxidative stress, and fat Peroxide in carcinogenesis and its contribution To start and develop the process well Documented in recent years, Using MDA as a sign of oxidative stress, there It was a growing interest in studying the role it played Lipid peroxide in the development of cancer. MDA low molecules Weight of aldehyde that can be produced free of charge Radical attack on
- Akbulut, H., Akbulut, K. G., Icli, F., & Büyükcelik, A. (2003). Daily variations of plasma malondialdehyde levels in patients with early breast cancer. *Cancer Detection and Prevention, 27*(2), 122-126.
- Basu, S., Harris, H., Wolk, A., Rossary, A., Caldefie-Chézet, F., Vasson, M.-P., & Larsson, A. (2016). Inflammatory F2-isoprostane, prostaglandin F2 $\alpha$ , pentraxin 3 levels and breast cancer risk: The Swedish Mammography Cohort. *Prostaglandins, Leukotrienes and Essential Fatty Acids, 113*, 28-32.
- Bennett, A., Stamford, I., Berstock, D., Dische, F., Singh, L., & A'Hern, R. (1989). Breast cancer, prostaglandins and patient survival. *British journal of cancer, 59*(2), 268.
- Dai, Q., & Zhu, X. (2009). F2-isoprostanes and metabolite, and breast cancer risk. *North American journal of medicine & science, 2*(3), 106.
- DeSantis, C. E., Bray, F., Ferlay, J., Lortet-Tieulent, J., Anderson, B. O., & Jemal, A. (2015). International variation in female breast cancer incidence and mortality rates. *Cancer Epidemiology and Prevention Biomarkers, 24*(10), 1495-1506.

- Israyelyan, A. H. (2003). The development of molecular diagnostics for breast cancer.
- Jabir, F.A. & Hoidy H. W. (2017). No Evaluation of Serum P53 Levels in Iraqi Female Breast Cancer Patients. *Asian Pacific Journal of Cancer Prevention.*, 18(9). 2551-2553
- Jabir, F.A. & Hoidy H. W. (2018). Pharmacogenetics as Personalized Medicine: Association Investigation of SOD2 rs4880, CYP2C19 rs4244285, and FCGR2A rs1801274 Polymorphisms in a Breast Cancer Population in Iraqi Women. *Clinical Breast Cancer*, 18(5), 863-868.
- Kang, D.-H. (2002). Oxidative stress, DNA damage, and breast cancer. *AACN Advanced Critical Care*, 13(4), 540-549.
- Lee, J. D., Cai, Q., Shu, X. O., & Nechuta, S. J. (2017). The role of biomarkers of oxidative stress in breast cancer risk and prognosis: A systematic review of the epidemiologic literature. *Journal of Women's Health*, 26(5), 467-482.
- Mannello, F., Tonti, G. A., Pagliarani, S., Benedetti, S., Canestrari, F., Zhu, W., . . . Sauter, E. R. (2007). The 8-epimer of prostaglandin F2 $\alpha$ , a marker of lipid peroxidation and oxidative stress, is decreased in the nipple aspirate fluid of women with breast cancer. *International Journal of Cancer*, 120(9), 1971-1976.
- Noda, N., & Wakasugi, H. (2001). Cancer and oxidative stress. *Japan Medical Association Journal*, 44(12), 535-539.
- Nourazarian, A. R., Kangari, P., & Salmaninejad, A. (2014). Roles of oxidative stress in the development and progression of breast cancer. *Asian Pac J Cancer Prev*, 15(12), 4745-4751.
- Pande, D., Negi, R., Khanna, S., Khanna, R., & Khanna, H. D. (2011). Vascular endothelial growth factor levels in relation to oxidative damage and antioxidant status in patients with breast cancer. *Journal of breast cancer*, 14(3), 181-184.
- Sahu, A., Varma, M., & Kachhawa, K. (2015). A prognostic study of MDA, SOD and catalase in breast Cancer patients. *Int J Sci Res*, 4(5), 157-159.
- Tillashaykhov, M., Khudaykulov, A., Danilova, E., Usupbekov, A., & Alimkhodjaeva, L. (2017). Study of trace elements status of breast cancer patients. *Онкологія*.